CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Astro 250 EC Growth Regulator

ACTIVE CONSTITUENT: 250 g/L TRINEXAPAC-ETHYL

For the Reduction of Leaf and Stem Growth of Grass Species, as an Aid in Turf Management, and as an Aid in Winter Grass Management

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA Approval No.: 64369/131605_B



Turf Culture 43 Gap Road Sunbury Vic 3429 Ph: 1300 11 8873 (Ph: 1300 11 TURF) www.turfculture.com.au

16-12-2022

Restraints

DO NOT use on bowling greens.

DO NOT apply to turf recovering from wear, renovation or turf affected by disease, waterlogging or other stresses.

DO NOT use where growth regulatory effects are likely following the application of a fungicide or any other material.

DO NOT apply to turf going into or coming out of dormancy.

DO NOT apply if rainfall is expected with 1 hour, and **DO NOT** irrigate for at least 1 hour after application.

DO NOT apply to Kentucky Bluegrass as damage may occur.

	Grasses	Rate		
Situation		Low Volume Spray /ha	Knapsack Application per 100 m ²	Critical Comments
Golf course greens	Bent grass	500 mL to 1 L	5 to 10 mL	 Apply Astro 250 EC Growth Regulator at least 6 hours before or after mowing and when turf is actively growing. Use the lower rates on turf with moderate growth rates where a lower level of growth reduction is required or when used in a program. Apply rates towards the higher end of the rate range when growth is vigorous or when longer periods of up to 6 weeks suppression is required. Application made to turf growing under cool conditions or to turf with low nutrient levels or where growth rates are low may result in discolouration. Well-maintained lawns and turf surfaces are those that receive at least weekly mowing and good levels of water and fertility are maintained. Refer to Application section for detailed information.
Golf course fairways Golf course short roughs, well- maintained lawns and turf surfaces	Common Couch	400 to 800 mL	4 to 8 mL	
	Couch varieties: Greenlees Park, Wintergreen, Santa ana	300 to 600 mL	3 to 6 mL	
	Fescue, Perennial Ryegrass	800 mL to 1.5 L	8 to 15 mL	
	Kikuyu, Carpet grass	800 mL to 2 L	8 to 20 mL	
	Buffalo grass, Common Couch, Hybrid Couch	500 mL to 1 L	5 to 10 mL	
	Fescue, Perennial Ryegrass, Bent grass	1 to 2 L	10 to 20 mL	
	Kikuyu	800 mL to 3 L	8 to 30 mL	
Golf course long roughs, parks, ovals, lawns, general grassed areas	Buffalo grass, Common Couch, Hybrid Couch	500 mL to 2 L	5 to 20 mL	
	Fescue, Perennial Ryegrass, Bent grass	1 to 3 L	10 to 30 mL	
	Kikuyu	1 to 4 L	10 to 40 mL	

	Grasses	Rate		
Situation		Low Volume Spray /ha	Knapsack Application per 100 m ²	Critical Comments
Bunker facing, path and garden edging, footpath and curbside, fencelines and around trees	Buffalo grass	2 L 20 mL	Apply Astro 250 EC Growth Regulator at least 6 hours before	
	Couch varieties: Greenlees Park, Wintergreen, Santa ana			or after mowing and when turf is actively growing.
				On difficult to mow areas apply Astro 250 EC Growth Regulator to actively growing turf and mow 3 to
	Common Couch, Perennial Ryegrass, Bent grass	2 to 3 L	20 to 30 mL	4 days after application. Apply in a strip about 25 to 50 cm wide.
				Use the lowest rate of Astro 250
	Kikuyu	3 to 4 L	30 to 40 mL	EC Growth Regulator around bunker edges and the highest rates in the other situations.
				Application made to turf growing under cool conditions or to turf with low nutrient levels or where growth rates are low may result in discolouration.

Turf Growth Management (Growth Regulation) continued

Winter Grass (Poa annua) Management

		Rate		
Situation	Grasses	Low Volume Spray/ha	Knapsack Application/ 100 m ²	Critical Comments
Golf course fairways or Bent grass golf greens	Winter Grass (<i>Poa annua</i>)	200 to 500 mL	2 to 5 mL	See instructions under Winter Grass Management for reduction of Winter Grass swards in Bent Grass golf greens. See Critical Comments aligned with Golf course greens situation/use for application recommendations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD

DO NOT graze treated turf/lawn, or feed turf/lawn clippings from any treated area to poultry or livestock.

GENERAL INSTRUCTIONS

Astro 250 EC Growth Regulator should be applied to actively growing turf, at least 6 hours before or after mowing. Good mowing management leading up to application is required to achieve a good quality turf surface and to obtain the best results from Astro 250 EC Growth Regulator.

Turf Growth Management (Growth Regulation)

The recommended use rates are designed to give approximately a 20 to 50% reduction in clippings over a 2 to 6 week period. Full growth regulation by Astro 250 EC Growth Regulator begins about 3 to 5 days after application. In practise the amount of growth reduction achieved will vary between situations due to environmental conditions and management practices. To some extent therefore application rates will need to be adjusted to match growing conditions, management practises and the amount of growth regulation required by the turf manager.

Repeat applications can be made as soon as turf resumes growth or more suppression is required.

Areas treated with Astro 250 EC Growth Regulator should continue to receive regular maintenance practises, including irrigation, fertilisation, and weed, disease and insect control where necessary, as recommended for quality turf.

Winter Grass (Poa annua) Management

Different grasses vary in their sensitivity to Astro 250 EC Growth Regulator. Since Winter Grass is more sensitive to Astro 250 EC Growth Regulator than Bent grass, Astro 250 EC Growth Regulator can be used to reduce the competitiveness of Winter Grass in Bent grass golf greens. Astro 250 EC Growth Regulator can be integrated into existing Winter Grass control programs. Low rates of Astro 250 EC Growth Regulator (that provide growth suppression of Winter Grass, but not Bent grass) applied during peak growth periods may enable a slow transition in the golf green sward composition away from Winter Grass and towards higher percentage of Bent grass.

Mixing

Add the required quantity of Astro 250 EC Growth Regulator directly to a spray tank containing 2/3 of the required spray volume. Add the rest of the water and ensure the mix is thoroughly agitated before application.

Application

Ensure even application in order to achieve even growth reduction. Avoid overlapping sprays.

Application Low Volume: Apply in 100 to 400 L/ha water using suitable flat fan nozzles. Application volume should be adequate to ensure thorough and even coverage of the turf leaves and penetration to the crowns. In higher cut turf (>15 cm) a significant spray shielding effect can occur, impacting negatively on spray penetration and even coverage at low application volumes.

Knapsack: It is important to apply the correct quantity of Astro 250 EC Growth Regulator to a known area of turf. The water volume used is not as important providing good coverage is achieved. Apply between 2 and 10 L of water per 100 m² of turf. Preferably use flat fan nozzles and ensure even coverage is achieved.

<u>Golf Course Greens</u>: Ensure product placement as uniformly as possible onto leaves and into crowns. Ideal application volume should be 350 to 500 L/ha. For best results use medium [XR Teejet* (11004 or 11005)] to coarse [Turbo Teejet* or AIXR Teejet* (11004 or 11005)] nozzles, at 5 km/hr and 3 bar pressure. Boom height should not exceed 50 cm above the ground.

<u>All Other Turf Areas</u>: Ensure product placement as uniformly as possible onto leaves and into crowns. Ideal application volume should be 200 to 500 L/ha. For best results use medium [XR Teejet* (11004 to 11008)] to coarse [Turbo Teejet* or AIXR Teejet* (11004 or 11005)] nozzles, at 8 km/hr and 3 bar pressure. Boom height should not exceed 50 cm above the ground.

Compatibility

Astro 250 EC Growth Regulator is compatible with Azoxy 95 Fungicide and Thumper Insecticide. For Common Couch, Hybrid Couch, and Kikuyu turf grasses only:

Astro 250 EC Growth Regulator is compatible with some formulations of products containing the following chemicals:

MCPA ester/bromoxynil, MCPA/bromoxynil/dicamba, bromoxynil, MCPA ester/bromoxynil/dicamba, MCPP/2,4-D amine/dicamba, 2,4-D amine, MCPP/2,4-D amine/dicamba or MCPP/dicamba. However, some transient discolouration of the treated turf may occur, after application of a tank mix. For the treatment of broadleaf weeds in turf with herbicides other than those indicated above, it is recommended to apply Astro 250 EC Growth Regulator and the broadleaf weed herbicides separately, allowing at least 10 days between an application of Astro 250 EC Growth Regulator and the broadleaf weed herbicides.

PRECAUTIONS

Re-entry Period

DO NOT enter treated areas, without protective clothing until spray has dried. DO NOT allow the public to enter treated areas until spray has dried.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to fish larvae and oysters. DO NOT contaminate dams, rivers, streams or storm water drains with concentrate or used container. Wash out spraying equipment, mixing and measuring utensils with water onto grass.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product. **For refillable containers**

Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

SAFETY DIRECTIONS

Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and preparing spray wear cotton overalls, buttoned to the neck and wrist and elbow-length PVC gloves. After each day's use, wash gloves, and contaminated clothing. Wash hands after use.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which is available from the supplier or from the manufacturer's website: www.turfculture.com.au

CONDITIONS OF SALE

The use of Astro 250 EC Growth Regulator being beyond the control of the manufacturer, no warranty expressed or implied is given by Turf Culture regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Turf Culture accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

* Trademark

In a Transport Emergency Dial 000 Police or Fire Brigade