


CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Columbus Insecticide

ACTIVE CONSTITUENT: 250 g/L THIAMETHOXAM

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

**For Control of African Black Beetle, Argentinian Scarab
and Billbug in Turf**

IMPORTANT: READ THIS LEAFLET BEFORE USING THIS PRODUCT

APVMA Approval No.: 68967/134137



Turf Culture
43 Gap Road
Sunbury Vic 3429
Ph: 1300 11 8873
(Ph: 1300 11 TURF)
www.turfculture.com.au

DIRECTIONS FOR USE**Restraints****DO NOT** apply aerially.**DO NOT** apply if heavy rain is forecast.

Situation	Pests	Rate	Critical Comments
Turf, lawns	1st instar larvae of African Black Beetle (<i>Heteronychus arator</i>)	800 mL to 1.2 L per ha in at least 500 L water per ha	Apply prior to or at peak egg hatch, mid spring to early summer, or when small larvae are present. Ensure product placement as close to soil surface as possible. Total application volume should not be lower than 500 L per ha (5 L per 100 m ²). For best results use extremely coarse droplets [Turbo Floodjet* (TF5) or TurfJet* (TTJ10)]. Preferably spray onto wet or dewy grass. Irrigate with 6 to 12 mm of water commencing within 1 hour of application.
	1st and 2nd instar larvae of Argentinian Scarab (<i>Cyclocephala signaticollis</i>)	(8 to 12 mL per 100 m ² in at least 5 L water per 100 m ²)	
	Larvae of Billbug (<i>Sphenophorus brunnipennis</i>)		Monitor adult activity through late spring and early summer. Spray when numbers peak or when small larvae (4 mm) are found in the thatch or surface soil. Early application is essential to minimise grass damage due to feeding. Use the higher rate when pest pressure is high or 2nd instar larvae are present. Ensure product placement as close to soil surface as possible. Total application volume should not be lower than 500 L per ha (5 L per 100 m ²). For best results use extremely coarse droplets [Turbo Floodjet* (TF5) or TurfJet* (TTJ10)]. Preferably spray onto wet or dewy grass. Irrigate with 6 to 12 mm of water commencing within 1 hour of application.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

WITHHOLDING PERIOD**DO NOT** graze treated turf/lawn or feed turf/lawn clippings from any treated area to poultry or livestock.**GENERAL INSTRUCTIONS****Application**

To be effective Columbus Insecticide requires to be placed as close to the soil surface as possible. Ensure that equipment is properly calibrated to give an even distribution at the correct volume. Apply in a minimum of 500 L water per ha (5 L per 100 m²). For best results use extremely coarse droplets [Turbo Floodjet* (TF5) or TurfJet* (TTJ10)].

Mixing

Add the required amount of Columbus Insecticide to clean water in half-filled spray tank with the agitator or by-pass in operation. Maintain agitation while filling tank with remainder of water. Agitate whilst spraying.

Compatibility

As formulations of other manufacturer's products are beyond the control of Turf Culture, and water quality varies with location, all mixtures should be tested prior to mixing commercial quantities.

INSECTICIDE RESISTANCE WARNING

GROUP	4A	INSECTICIDE
--------------	-----------	--------------------

For insecticide resistance management Columbus Insecticide is a Group 4A insecticide. Some

naturally occurring insect biotypes resistant to Columbus Insecticide and other Group 4A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Columbus Insecticide or other Group 4A insecticides are used repeatedly. The effectiveness of Columbus Insecticide on resistant individuals could be significantly reduced.

Since occurrence of resistant individuals is difficult to detect prior to use, Turf Culture accepts no liability for any losses that may result from failure of Columbus Insecticide to control resistant insects. Columbus Insecticide may be subject to specific resistance management strategies. For further information contact your supplier, Turf Culture representative or local agricultural department agronomist.

PRECAUTIONS

Re-entry Period

DO NOT enter treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

PROTECTION OF LIVESTOCK

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. DO NOT graze treated turf/lawn, or feed turf/lawn clippings from any treated area to poultry or livestock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Toxic to certain aquatic invertebrates. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray drift onto adjacent areas, particularly wetlands, water-bodies or watercourses. DO NOT contaminate streams, rivers or waterways with the product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight. Triple-rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When opening the container, mixing and loading, preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. If applying by spraying equipment carried on the back of the user, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use and after each day's use, wash gloves and contaminated clothing.

FIRST AID INSTRUCTIONS

If poisoning occurs contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet (SDS), which is available from the supplier or from the manufacturer's website: www.turfculture.com.au

CONDITIONS OF SALE

The use of Columbus Insecticide being beyond the control of the manufacturer, no warranty expressed or implied is given by Turf Culture regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Turf Culture accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

* Trademark

In a Transport Emergency Dial 000 Police or Fire Brigade